

## LA-UR-21-24249

Approved for public release; distribution is unlimited.

Title: MUSiC: Measurement of Uranium Subcritical and Critical (IER 488)

Author(s): Weldon, Robert Allen Jr.  
Cutler, Theresa Elizabeth  
Hutchinson, Jesson D.  
Myers, William L.  
McKenzie, George Espy IV  
McSpaden, Alexander Thomas  
Sanchez, Rene Gerardo


Intended for: Web

Issued: 2021-05-03

---

**Disclaimer:**

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.



Delivering science and technology  
to protect our nation  
and promote world stability

# MUSiC: Measurement of Uranium Subcritical and Critical (IER 488)

Robert Allen Weldon Jr, Theresa Elizabeth  
Cutler, Jesson D. Hutchinson, William L.  
Myers, Geordie Espy McKenzie IV,  
Alexander Thomas McSpaden, and Rene  
Gerardo Sanchez



Nuclear Engineering and  
Nonproliferation Advanced  
Nuclear Technology (NEN-2)

[raw@lanl.gov](mailto:raw@lanl.gov)


2/24/2021





# Measurement of Uranium Subcritical and Critical (MUSiC)

- Measurements of bare, HEU system
  - Rocky Flats shells – variable mass
  - Ten configurations will be measured, spanning from deeply subcritical to critical (i.e., it is a measurement of similar systems with different reactivities)
- Key benefits of MUSiC:
  - Large range of reactivities (subcrit to crit)
    - First measurement of a simple/fast system that covers such a range to be submitted to ICSBEP
    - Subcritical benchmarks – useful for testing correlated fission physics (CGMF, FREYA)
  - Simple HEU system (new Lady Godiva HMF001) – valuable for nuclear data validation
  - Multiple detector systems and measurement methods
    - Provide results independent of the detector system
    - Demonstrate the suitability of organic scintillators for future benchmark measurements




# Rocky Flats HEU hemisphere shells

- 93.16% U-235
- ~0.3 cm thick
- Variable radius between ~2 cm and 10 cm
- MUSiC configurations:



Simulated configurations

Configuration	Mass (kg)	$k_{eff}$
1	15.723	0.64925
2	21.832	0.72715
3	33.552	0.83854
4	38.243	0.87888
5	47.925	0.92428
6	54.642	0.97914
7	60.190	0.99179
8	60.657	1.00228
9	61.009	1.01077
10	67.082	1.02448




# Measurement details

- Induced fission driven by:
  - DT generator
  - Cf-252
- Detectors
  - NoMAD
    - 15 He-3 tubes embedded in high density polyethylene
    - Portable
  - Array of 4 small volume He-3 detectors
    - High pressure (40 atm vs 10 atm for NoMAD)
    - Fast recovery speed – suitable for Rossi- $\alpha$  measurements
  - **Array of 8, 2"×2" EJ-309 detectors**
    - New detection system for NCERC
    - Direct detection of fast neutrons
    - No moderation, fast response → ideal for bare systems



# Approach to critical (12/20)

- Measurements to determine the final configurations were performed in December 2020
  - Approach to critical
  - Remeasured shell masses (losses)
- Changes between simulated and experimentally determined configurations:



Experimentally determined configurations

Configuration	Mass (kg)	$k_{eff}$
1	13.0428	0.64
2	21.6432	0.74
3	29.0415	0.81
4	37.9617	0.88
5	42.9722	0.91
6	48.4099	0.95
7	54.2785	0.98

Simulated configurations

Configuration	Mass (kg)	$k_{eff}$
1	15.723	0.64925
2	21.832	0.72715
3	33.552	0.83854
4	38.243	0.87888
5	47.925	0.92428
6	54.642	0.97914
7	60.190	0.99179
8	60.657	1.00228
9	61.009	1.01077
10	67.082	1.02448

# The Rossi- $\alpha$ method

- Goal: infer reactivity from a measurement of the prompt neutron decay
  - Prompt neutron decay constant ( $\alpha$ ) – rate of prompt neutron population change with time (units of inverse time)
  - Represents an eigenvalue characteristic of a system that can (in principle) be calculated and measured
- Assumptions in relation to reactivity
  - Based on point kinetic reactor model
  - The spatial flux is time independent
  - Not true for subcritical systems, but it is a reasonable approximation down to some (?) subcritical value of  $k_{eff}$
- Method compliments Feynman variance-to-mean and pulsed source techniques

Reactivity and  $k_{eff}$ :

$$\rho = \frac{k_{eff} - 1}{k_{eff}}$$

Reactivity and  $\alpha$ :

$$\rho = \beta_{eff} - \alpha\Lambda$$

$\beta_{eff}$  – effective  
delayed neutron  
fraction


$\Lambda$  – mean neutron  
generation time

# Implementation of the Rossi- $\alpha$ method


- Simple measurement
  - Collect neutron interaction times using the detectors
  - Analyze events within a selected time window (or reset time)
  - Calculate time difference between interactions (type 1 binning)
  - Number of counts will be correlated to the time difference
  - Estimate  $\alpha$  by fitting the curve with

$$p(t) = Ae^{-\alpha t} + B \quad [1]$$

- Accurate for the time dependence of the prompt neutron decay for a fissioning assembly at low power (need to avoid overlapping fission chains)



Binning types. Figure from [1].






[1] Hansen, Gordon E. *Rossi Alpha Method*. No. LA-UR-85-4176; CONF-8508105-7. Los Alamos National Lab., NM (USA), 1985.



# EJ-309 liquid organic scintillators

- Time resolution ( $\sim 900$  ps)
  - Excellent for neutron noise measurements
- Pulse shape discrimination – eliminate gamma interactions
- Energy information is also available – spectroscopy





From [2].

[2] Brown, J. A., Goldblum, B. L., Laplace, T. A., Harrig, K. P., Bernstein, L. A., Bleuel, D. L., & Marleau, P. (2018). Proton light yield in organic scintillators using a double time-of-flight technique. *Journal of Applied Physics*, 124(4), 045101.

# Conclusions and future work

- MUSiC
  - Bare, HEU measurements covering a large range of reactivities
  - Multiple detector systems and analysis methods are being used/implemented to obtain detector-independent results
- Future work
  - Measurements: February – March 2021
  - Simulations
    - Event-by-event simulations of Rossi- $\alpha$
    - ACODE – under development for MCNP6
    - MCATK – can be used to calculate subcritical  $\alpha$ -eigenvalues





# Acknowledgements

- This work was supported by the DOE Nuclear Criticality Safety Program, funded and managed by the National Nuclear Security Administration for the Department of Energy



# Simulations (KCODE, MCNPX-PoliMi)


- MCNP6 KCODE

- Calculates  $k_{eff}$
- KOPTS card used to calculate  $\beta_{eff}$  and  $\Lambda$
- Reactivity and  $\alpha$ :

$$\rho = \frac{k_{eff} - 1}{k_{eff}} = \beta_{eff} - \alpha\Lambda$$


- MCNPX-PoliMi

- Simulations of Rossi- $\alpha$  measurement
  - Same geometry as KCODE
  - Analysis is performed on event-by-event neutron interactions in the EJ-309 detectors
- Fit to estimate  $\alpha$
- Use  $\beta_{eff}, \Lambda$  from KOPTS



# Rossi- $\alpha$ simulated results

- Good agreement between KCODE and PoliMi results
- Uncertainty in  $k_{eff}$  estimated with PoliMi is mostly attributed to the uncertainty in  $\Lambda$  calculated with KCODE
- Comparison to GODIVA IV measurements (also bare HEU)
  - $k_{eff} = 0.995$
  - $\alpha^{-1} = 1180 \text{ ns}$



Configuration	MCNP KCODE				MCNPX-PoliMi				$\chi^2$
	$\alpha^{-1}$ (ns)	$\sigma_{\alpha^{-1}}$ (ns)	$k_{eff}$	$\sigma_{k_{eff}}$	$\alpha^{-1}$ (ns)	$\sigma_{\alpha^{-1}}$ (ns)	$k_{eff}$	$\sigma_{k_{eff}}$	
2	36	4	0.7289	$3 \times 10^{-5}$	38.3	0.4	0.72	0.02	1.8
3	85	9	0.84455	$4 \times 10^{-5}$	58.3	0.7	0.84	0.02	1.19
4	81	6	0.88034	$4 \times 10^{-5}$	67.8	0.5	0.870	0.009	1.27
5	170	20	0.92745	$4 \times 10^{-5}$	103.2	0.4	0.92	0.01	1.12
6	400	30	0.97956	$4 \times 10^{-5}$	322.2	0.5	0.977	0.002	1.05
7	880	80	0.99432	$4 \times 10^{-5}$	1177.2	0.6	0.9935	0.0008	1.01

# Deviation from fit with decreasing effective multiplication

- Rossi- $\alpha$  method is only applicable over regions of reactivity where the point kinetic model assumptions are true

$$p(t) = Ae^{-\alpha t} + B \quad [1]$$

Configuration	$\chi^2$
2	1.8
3	1.19
4	1.27
5	1.12
6	1.05
7	1.01

